#### Convex Optimization Part I

CS 584: Big Data Analytics

Material adapted from John Duchi (<u>https://www.cs.berkeley.edu/~jordan/courses/294-fall09/lectures/optimization/slides.pdf</u>) & Stephen Boyd (<u>https://web.stanford.edu/class/ee364a</u>)

### **Optimization Problem**

Minimize a function subject to some constraints

$$\min_{x} f_{0}(x)$$
  
s.t.  $f_{k}(x) \leq 0, k = 1, 2, \cdots, K$   
 $h_{j}(x) = 0, j = 1, 2, \cdots, J$ 

• Example: Minimize the variance of your returns while earning at least \$100 in the stock market.

# Machine Learning and Optimization

Linear regression

$$\min_{w} ||Xw - y||^2$$

Logistic regression r

$$\min_{w} \sum_{i} \log(1 + \exp(-y_i x_i^{\top} w))$$

• SVM 
$$\min_{w} ||w||^2 + C \sum_{i} \xi_i$$

s.t. 
$$\xi_i \ge 1 - y_i x_i^\top w$$
  
 $\xi_i \ge 0$ 

• And many more ...

#### Non-Convex Problems are Everywhere

- Local (non-global) minima
- All kinds of constraints



No easy solution for these problems



Consider convex problems

#### Convex Sets

A set C is convex such that given any two points a, b in that set, the line segment between the two points is in the set

 $x_1, x_2 \in C, 0 \le \theta \le 1 \implies \theta x_1 + (1 - \theta) x_2 \in C$ 



convex

concave

CS 584 [Spring 2016] - Ho

#### Examples: Convex Set

- Real space:  $\mathbb{R}^n$
- Non-negative orthant:  $\mathbb{R}^n_+$
- Norm balls:  $\{x \mid ||x x_c|| \le r\}$
- Hyperplane:  $\{x \mid a^{\top}x = b\}, a \neq 0$
- Halfspace:  $\{x \mid a^{\top}x \leq b\}, a \neq 0$

## Convexity-preserving operations

- Intersection
- Affine functions  $f(x) = Ax + b, A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m$ (e.g., scaling, translation, projection)
- Perspective function

$$P(x,t) = \frac{x}{t}, \text{ dom } P = \{(x,t) \mid t > 0\}$$

Linear-fractional functions

$$f(x) = \frac{Ax + b}{c^{\top}x + d}, \text{ dom } f = \{x \mid c^{\top}x + d > 0\}$$

#### **Convex Functions**

#### Definition

 $f: \mathbb{R}^n \to \mathbb{R}$  is convex if **dom** f is a convex set and  $f(\theta x + (1 - \theta)y) \le \theta f(x) + (1 - \theta)f(y)$ for all  $x, y \in$ **dom**  $f, 0 \le \theta \le 1$ 



CS 584 [Spring 2016] - Ho

#### Examples: Convex Functions (Real space)

- affine: ax + b, for any  $a, b \in \mathbb{R}$
- exponential:  $e^{ax}$ , for any  $a \in \mathbb{R}$
- powers:  $x^{\alpha}$ , for  $\alpha \ge 1$  or  $\alpha \le 0$ ,  $x \in \mathbb{R}_{++}$
- powers of absolute value:  $|x|^p$ , for  $p \ge 1$
- negative entropy:  $x \log x, x \in \mathbb{R}_{++}$

# Convex Optimization Problem

Definition:

An optimization problem is **convex** if its objective is a convex function, the inequality constraints are convex, and the equality constraints are affine

$$\begin{split} \min_{x} \ f_0(x) & \text{convex function} \\ \text{s.t.} \ f_k(x) \leq 0, k = 1, 2, \cdots, K & \text{convex sets} \\ h_j(x) = 0, j = 1, 2, \cdots, J & \text{affine constraints} \end{split}$$

# Benefits of Convexity

- Theorem: If x is a local minimizer of a convex optimization problem, it is a **global** minimizer
- Theorem: If the gradient at c is zero, then c is the global minimum of f(x)

$$\nabla f(c) = 0 \iff c = x^*$$

# Lagrange Duality

- Bound or solve an optimization problem via a different optimization problem
- Reformulate the problem as an augmented objective with a weighted sum of constraints
  - Remove constraints
  - Introduce new variables
  - Form a dual function

#### Constructing the dual

Original optimization problem

min  $f_0(x)$  $\boldsymbol{T}$ s.t.  $f_k(x) \le 0, k = 1, 2, \cdots, K$  $h_j(x) = 0, j = 1, 2, \cdots, J$ dual function  $g(\lambda, v) = \inf_{x} \left\{ f_0(x) + \sum_{k} \lambda_k f_k(x) + \sum_{i} v_j h_j(x) \right\}$  $\lambda_i \geq 0, v_i \in \mathbb{R}$ 

#### Two Properties of Dual

- Weak Duality (Lemma): If  $\lambda \ge 0$ , then  $g(\lambda, v) \le f_0(x^*)$ 
  - Always holds for convex and non convex problems
  - Can be used to find nontrivial lower bound for difficult problems
  - **Strong Duality** (Theorem):  $d^* = x^*$

•

- (Usually) holds for convex problems
- Constraint qualifications are conditions that guarantee strong duality in convex problems

# Unconstrained Optimization Algorithms



CS 584 [Spring 2016] - Ho

#### Gradient Descent (Steepest Descent)

- Simplest and extremely popular
- Main Idea: take a step proportional to the negative of the gradient
- Easy to implement
- Each iteration is relatively cheap
- Can be slow to converge



#### Gradient Descent Algorithm

Algorithm 1: Gradient Descent

while Not Converged do  

$$| x^{(k+1)} = x^{(k)} - \eta^{(k)} \nabla f(x)$$
  
end  
return  $x^{(k+1)}$ 



CS 584 [Spring 2016] - Ho

#### Importance of Step Size

Challenge is to find a good step size to avoid step size that is too long or too short



too long => divergence too short => slow convergence

#### Step Size Selection

• Exact Line Search: Pick step size to minimize the function

$$\eta^{(k)} = \argmin_{\eta} f(x - \eta \nabla f(x))$$
 Too expensive to be practical

 Backtracking (Armijo) Line Search: Iteratively shrink the step size until a decrease in objective is observed

> Algorithm 1: Backtracking Line Search Let  $\alpha \in (0, \frac{1}{2}), \beta \in (0, 1)$ while  $f(x - \eta \nabla f(x)) > f(x) - \alpha \eta ||\nabla f(x)||^2$  do  $| \eta = \beta \eta$ end

#### Example: Linear Regression

• Optimization problem:

$$\min_{w} ||Xw - y||_2$$

• Closed form solution:

$$w^* = (X^\top X)^{-1} X^\top y$$

• Gradient update:

$$w^{+} = w - \frac{1}{m} \sum_{i} (x_{i}^{\top} w - y_{i}) x_{i}$$

#### Some Resources for Convex Optimization

- Boyd & Landenberghe's Book on Convex Optimization <u>https://web.stanford.edu/~boyd/cvxbook/bv\_cvxbook.pdf</u>
- Stephen Boyd's Class at Stanford <u>http://stanford.edu/class/ee364a/</u>
- Vandenberghe's Class at UCLA <u>http://www.seas.ucla.edu/~vandenbe/ee236b/ee236b.html</u>
- Ben-Tai & Nemirovski Lectures on Modern Convex Optimization <u>http://epubs.siam.org/doi/book/10.1137/1.9780898718829</u>